DIRECTIONS FOR USE

When mixed, fill spray tank half full with water and add the required amount of Organic-N while continuing to fill tank with water. Continue agitation through the spray operation. See Precautions. Always agitate drum contents before use.

APPLICATION RATES

Horticultural Crops:

Minimum dilution 1:50 with water.

Foliar Spray: 2 - 5L/ha; repeat as required.

Irrigation: 5 - 10L/ha; repeat as required.

Broad-acre Crops and Pastures:

Minimum dilution 1:50 with water.

Foliar Spray: 2-5L/ha; repeat as required.

Each farmer's situation and processes and practices may differ and therefore necessitate corrections to ensure optimum results. Adjust rates according to production levels. Repeat as required.

Always test for compatibility and crop suitability prior to use. Seek advice for rates suited to your crop.

TYPICAL ANALYSIS

Nitrogen 10% w/v as amino acids. With fulvic acid 23% w/v, trace elements (Ca, Mg, Zn, Fe), Organic Carbon 37.5% w/v.

CLEAN UP PROCEDURE

Use all mixture in spray and irrigation tanks; purge tanks and lines with clean water; flush irrigation lines. Do not return mix to original drums.

ORGANIC FERTILISER - LIQUID

Concentrated Liquid N - 10%

- For managing crop nitrogen demand and building yield
- Provides immediate nutrition for your crop Ideal to stimulate plant vegetative growth
- With 23% fulvic acid to improve nutrient uptake
- 100% soluble liquid
- Use via irrigation or foliar spray
- Hydrolysed plant protein

200L 1,000L

20L

Shelf life: 3 years Manufacturing Date: Batch #







STORAGE

Store in original container away from direct sunlight. Do not store in diluted form. Do not store at temperatures exceeding 40°C.

PRECAUTIONS

Physically compatible with a wide range of commonly used products. Always mix a small quantity (jar test) and check for physical compatibility before combining with other ingredients.

When applying for the first time, or in combination with other products, a small test area should be sprayed and observed prior to the total spray.

Exceeding 1:50 concentration may create a phytotoxic reaction in some environmental conditions.

Caution should be used when applying as a foliar spray in conjunction with copper based products as increased activity of copper may produce unacceptable levels of phytotoxicity. Do not apply as foliar spray during flowering.

WARNING

Do not swallow. The spray from this product may act as a irritant. Avoid inhalation and contact with eyes and skin. Read SDS prior to use.

Wear protective clothing, safety glasses and gloves when using this product.

DISCLAIMER

Organic Farming Systems (OFS) has no control over farming methods. application of the product, irrigation systems, weather or growing conditions. Consequently no responsibility or liability for any failure in performance, losses, damages or injuries will be accepted by Sustainable Ventures Pty Ltd trading as OFS, its distributors or agents following the use of this product. Seek independent advice prior to use.



Organic Farming Systems Level 1, 322 Hay St, Subiaco 6008. Tel: 08 9388 3623 Tel: 03 9008 6352

info@organicfarming.com.au www.organicfarming.com.au

